



Visualization and Analysis of Near Real-Time Global Cloud Composites (GCC):

Integration in a Geospatial Web Mapping Application

Valeria Green

Booz Allen Hamilton Inc.

Co-Authors: Louis Nguyen 1, William L. Smith, Jr. 1, Thad L. Chee2, Audy J. Barnett3, Benjamin M. Landes3

¹NASA Langley Research Center, Hampton, Virginia

²ADNET Systems, Inc., Bethesda, Maryland

³Booz Allen Hamilton Inc., McLean, Virginia



Esri Cloud Raster Format



Multidimensional/Transposed Cloud-optimized analysis ready data (ARD)

NASA Earthdata GIS (gis.earthdata.nasa.gov)



Geospatially service enabled ArcGIS Image Services hosted at Earthdata GIS

Esri API for JS v4 & Calcite Design



Custom web mapping application with visualization analytical functionality

Technologies Used on Web App

React/JavaScript

JSX

CSS

React Redux
ArcGIS Maps SDK for JavaScript
Calcite Design System



Imagery Analysis Capabilities

Select and Manipulate Imagery

Toggle visibility and customize with a pixel filter, transparency control and color ramp table/slider

Compare Multiple Layers

Compare multiple layers with a swipe widget

Time slider

Animate temporally enabled imagery layers with a slider

Time series

Display a time series of the data from temporally enabled imagery layers





Field Campaigns Support

Search Functionality

Filter field campaigns by name

Detailed Field Campaign View

View details about each field campaign individually

Detailed Flight Calendar

View flight data for specific days within a calendar layout

Direct Access to Data Files

Download the direct data relevant to the field campaign





Direct Data Subset & Download

Subset vs Full

Subset the imagery data from OpenDAP

Change Subset Data Format

Change the format of the subset data which include PNG, KMZ, GeoTIFF and NetCDF

Change Subset Time

Select the time and date of the data to subset with

Populate from Rectangle Tool

Use the rectangle sketch tool to directly populate coordinates or vice versa



